

March 25, 2019, Monday

Which of these is NOT a function? → Vertical Line Test!

A. $(5, 3), (6, 4), (7, 3), (8, 4)$

B.

C. $y = 3x^2$

D.

Let's review factoring quadratics.....

Algebra 1 Name _____ ID: 1
Assignment Date _____ Period _____
Solve each equation by factoring.
1) $x^2 - 9x + 8 = 0$
 $(x-8)(x-1) = 0$
 $x-8 = 0 \quad x-1 = 0$
 $x = 8 \quad x = 1$

2) $2x^2 - x - 3 = 0$
 $2x^2 - x - 3 = 0$
 $(x+1)(2x-3) = 0$
 $x+1 = 0 \quad 2x-3 = 0$
 $x = -1 \quad x = \frac{3}{2}$

Solve each equation by taking square roots.
3) $x^2 = 500$
 $\sqrt{x^2} = \sqrt{500}$
 $x = \pm 10$
 $x = 10, x = -10$

Solve each equation by completing the square.
4) $p^2 + 6p + 9 = 0$
 $p^2 + 6p + 9 = 0$
 $(p+3)^2 = 0$
 $p+3 = 0$
 $p = -3$

$p+3 = \pm 3$
 $p = -3 \pm 3$
 $p = -3 + 3 = 0$
 $p = -3 - 3 = -6$

Algebra 2 Name _____
TODD Solving Quadratics (Factoring/Sq. Root/Completing the Square)

Solve each equation by factoring.
1) $a^2 - b^2 = (a+b)(a-b)$
2) $n^2 - 16 = 0$
 $a^2 - b^2 = (a+b)(a-b)$
 $(n+4)(n-4) = 0$
 $n+4 = 0 \quad n-4 = 0$
 $n = -4 \quad n = 4$

3) $x^2 + 9x + 14 = 0$
4) $5m^2 - 23m - 10 = 0$

Solve each equation by taking square roots.
5) $9x^2 = 9$
 $\sqrt{9x^2} = \sqrt{9}$
 $3x = \pm 3$
 $x = \pm 1$

Solve each equation by completing the square.
7) $x^2 - 4x + 3 = 0$
8) $k^2 + 12k + 35 = 0$

Algebra 2 Name _____
TODD Solving Quadratics (Factoring/Sq. Root/Completing the Square)

Solve each equation by factoring.
1) $n^2 - 21 = 0$
 $(n+3)(n-7) = 0$
 $n+3 = 0 \quad n-7 = 0$
 $n = -3 \quad n = 7$

2) $x^2 - 9 = 0$

Solve each equation by taking square roots.
5) $4x^2 + 2x - 401 = 0$
 $\sqrt{4x^2 + 2x - 401} = \sqrt{0}$
 $2x = \pm \sqrt{401}$
 $x = \pm \frac{\sqrt{401}}{2}$

6) $8x^2 - 5 = 387$
 $8x^2 = 392$
 $\sqrt{8x^2} = \sqrt{392}$
 $x = \pm \sqrt{\frac{392}{8}}$
 $x = \pm \sqrt{49}$
 $x = \pm 7$

Solve each equation by completing the square.
7) $x^2 + 16x + 26 = 0$
 $x^2 + 16x + 64 = 38$
 $(x+8)^2 = 38$
 $x+8 = \pm \sqrt{38}$
 $x = -8 \pm \sqrt{38}$
 $x = -8 + \sqrt{38} \approx -1.4$
 $x = -8 - \sqrt{38} \approx -17.4$

8) $n^2 - 16n + 60 = 0$

sg....

March 26, 2019, Tuesday

1. What is the y-intercept of the graph of $h(x) = 2^x - 4$?
A. $(0, -4)$
B. $(0, -3)$
C. $(0, 1)$
D. $(0, 2)$

2. What is the range of the graph of $f(x) = -3(x-4)$?
A. $(-3, 4)$
B. $(-3, \infty)$
C. $(-\infty, 4)$
D. $(-\infty, \infty)$

Answers to Unit 5.3 S
1. B 2. D

Algebra 1 ~ Day 4
Solving by the Quadratic Formula Notes
The solutions of any quadratic equation $(ax^2 + bx + c = 0)$ can be found by evaluating the quadratic formula:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Example: Use the quadratic formula to solve for x :

$$\begin{aligned} 1. & 2x^2 - 10x - 5 = 0 \\ & a = 2, b = -10, c = -5 \\ & x = \frac{-(-10) \pm \sqrt{(-10)^2 - 4(2)(-5)}}{2(2)} \\ & x = 5.5, -0.5 \end{aligned}$$

$$\begin{aligned} 2. & 9x^2 + 2 = 3x \\ & 9x^2 - 3x + 2 = 0 \\ & a = 9, b = -3, c = 2 \\ & x = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(9)(2)}}{2(9)} \\ & x = \text{NO SOLUTIONS} \end{aligned}$$

$$\begin{aligned} 3. & -x^2 - 6x = 9 \\ & x^2 + 6x + 9 = 0 \\ & a = 1, b = 6, c = 9 \\ & x = \frac{-6 \pm \sqrt{(-6)^2 - 4(-1)(9)}}{2(-1)} \\ & x = -3, -3 \end{aligned}$$

Algebra 1 ~ U3B Day 4
Solving by the Quadratic Formula
Solve each equation with the quadratic formula.

1) $4p^2 - 9 = 0$ 2) $6p^2 + 8p - 30 = 0$
 $a = 4, b = 0, c = -9$ $a = 6, b = 8, c = -30$
 $p = \pm \sqrt{\frac{9}{16}}$ $p = \frac{-8 \pm \sqrt{64 + 144}}{12}$
 $p = \pm \frac{3}{4}$ $p = \frac{-8 \pm 10}{12}$
 $p = \pm 0.75$ $p = \frac{1}{6}, -5$

3) $6b^2 + 7b - 68 = 0$ 4) $5x^2 + 11 = 0$
 $a = 6, b = 7, c = -68$ $a = 5, b = 0, c = 11$
 $b = \pm \sqrt{\frac{49 + 176}{36}}$ $x = \pm \sqrt{\frac{11}{25}}$
 $b = \pm \sqrt{\frac{225}{36}}$ $x = \pm \sqrt{\frac{11}{25}}$
 $b = \pm 5$ $x = \pm \frac{\sqrt{11}}{5}$
 $x = 2.833, -4$ $x = \pm 1.5$

5) $r^2 + 4r - 49 = -9$ 6) $2a^2 - 58 = -8$
 $r^2 + 4r + 49 = 40$ $a^2 - 29 = 4$
 $(r+7)^2 = 40$ $a = \pm \sqrt{33}$
 $r+7 = \pm \sqrt{40}$ $a = \pm \sqrt{33}$
 $r = -7 \pm \sqrt{40}$ $a = \pm \sqrt{33}$

7) $4t^2 - 8t = 21$ 8) $4b^2 = 144$
 $t^2 - 2t = \frac{21}{4}$ $b^2 = 36$
 $t^2 - 2t - 5.25 = 0$ $b = \pm 6$
 $(t-1)^2 = 6.25$ $b = \pm 6$
 $t-1 = \pm \sqrt{6.25}$ $b = \pm 6$
 $t = 1 \pm 2.5$ $b = \pm 6$

9) $2a^2 - 5 + 3a = -10 - 8a^2 + 3a$
 $10) m^2 + 8m = 91 + 2m$
 $m^2 + 6m - 91 = 0$ $a = 1$
 $m^2 + 6m - 91 = 0$ $b = 6$
 $m = -6 \pm \sqrt{36 + 364}$ $c = 91$
 $m = -6 \pm \sqrt{400}$
 $m = -6 \pm 20$
 $m = 14, -26$

11) $a^2 - 3a - 45 = 9a$
 $12) 12p^2 - 9p - 44 = -4p + 8p^2$

13) $2d^2 + 11d - 138 = 0$
A) $[6, -11.5]$
B) No solution.
C) $[2.231, -7.731]$
D) $[18.471, -7.471]$

14) $6n^2 + 7n - 20 = -7$
A) $[7, -7]$
B) $[6, -1.5]$
C) $[9.899, -9.899]$
D) $[1, -2.167]$
 $6n^2 + 7n - 13 = 0$
 $n = -7 \pm \sqrt{49 + 104}$
 $n = -7 \pm \sqrt{153}$
 $n = -7 \pm 12.36$
 $n = 5.36, -19.36$

15) $4d^2 = 24$
A) $[4, -4]$
B) $[2.5, -2.5]$
C) $[0.667, 0.5]$
D) $[1.25, -1.25]$

16) $6x^2 - x - 64 = 0$
A) $[2.833, -1.833]$
B) $[12.458, -5.458]$
C) $[1.198, -2.365]$
D) $[2, -6.5]$

Algebra 1
TODD Solving Quadratic Equations (any method)
Solve each equation your way (by factoring, square root, completing the square, or quadratic formula). SHOW ALL OF YOUR WORK!!!

1) $p^2 + 2p - 143 = 0$ 2) $4x^2 - 9 = 0$
 $(p+11)(p-13) = 0$ $2x = \pm 3$
 $p = -11, 13$ $x = \pm \frac{3}{2}$

3) $4x^2 - 7x - 15 = 0$ 4) $4v^2 + 3v - 27 = 0$
 $(4x+5)(x-3) = 0$ $4v^2 + 3v - 27 = 0$
 $x = -\frac{5}{4}, 3$ $v = -\frac{9}{4}, \frac{3}{4}$

Algebra 1
TODD Solving Quadratic Functions
Solve each equation your way (by factoring, square root, completing the square, or quadratic formula). SHOW ALL OF YOUR WORK!!!

1) $4k^2 - 81 = 0$ 2) $2k^2 - 4k - 16 = 0$
 $(2k+9)(2k-9) = 0$ $2k^2 - 4k - 16 = 0$
 $k = \pm 4.5$ $2k^2 - 4k - 16 = 0$
 $k = \pm 4$

3) $3b^2 - 4b - 7 = 0$ 4) $4v^2 - 2v - 6 = 0$
 $3b^2 - 4b - 7 = 0$ $4v^2 - 2v - 6 = 0$
 $(3b+7)(b-1) = 0$ $4v^2 - 2v - 6 = 0$
 $b = -\frac{7}{3}, 1$ $v = \frac{1}{2}, -\frac{3}{2}$

Algebra 1
TODD Solving Quadratic Functions
Solve each equation by factoring.

1) $x^2 + 5x + 6 = 0$ 2) $n^2 - n - 12 = 0$
 $(x+2)(x+3) = 0$ $(n+3)(n-4) = 0$
 $x = -2, -3$ $n = -3, 4$
 $x = -2, -3$ $n = -3, 4$
 $x = -2, -3$ $n = -3, 4$
 $x = -2, -3$ $n = -3, 4$

GCF: $3) 7r^2 + 3r = 0$
 $r(7r+3) = 0$
 $r=0$ $7r+3=0$
 $r=\frac{-3}{7}$

Solve each equation by taking the square root.

4) $a^2 = 1$
 $a = \pm 1$

5) $p^2 = 81$
 $p = \pm 9$

+ don't forget you can decimals or fractions

6) $64r^2 = 64$
 $64r^2 = 64$
 $r^2 = 1$
 $r = \pm 1$

Solve each equation by completing the square.

$$7) x^2 + 12x + 8 = 0$$

$$x^2 + 12x = -8$$

$$(x+6)^2 = 45$$

$$7(x+6)^2 = 45 + 36$$

$$7(x+6)^2 = 81$$

$$x+6 = \pm 9$$

$$x = -6 \pm 9$$

$$x = 3$$

$$x = -15$$

** You can have decimals or fractions!*

Solve each equation by completing the square formula.

$$8) \frac{9x^2}{2} - 10x + 11 = 0$$

$$\frac{9}{2}x^2 - 10x + 11 = 0$$

$$9x^2 - 20x + 22 = 0$$

$$a = 9, b = -20, c = 22$$

$$b^2 - 4ac = (-20)^2 - 4(9)(22)$$

$$b^2 - 4ac = 400 - 792$$

$$b^2 - 4ac = -392$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-(-20) \pm \sqrt{(-20)^2 - 4(9)(22)}}{2(9)}$$

$$x = \frac{20 \pm \sqrt{400 - 792}}{18}$$

$$x = \frac{20 \pm \sqrt{-392}}{18}$$

$$x = \frac{20 \pm \sqrt{392}}{18}$$

$$x = \frac{20 \pm 2\sqrt{98}}{18}$$

$$x = \frac{20 \pm 2\sqrt{49 \cdot 2}}{18}$$

$$x = \frac{20 \pm 2 \cdot 7\sqrt{2}}{18}$$

$$x = \frac{20 \pm 14\sqrt{2}}{18}$$

$$x = \frac{10 \pm 7\sqrt{2}}{9}$$

$$x = 1.6$$

Solve each equation by factoring, use any factoring method you choose.

$$9) x^2 + 7k + 6 = 0$$

$$X$$

$$GCF(1) \quad x^2 + 7x + 6 = 0$$

$$x^2 + 6x + x + 6 = 0$$

$$x(x + 6) + 1(x + 6) = 0$$

$$(x + 1)(x + 6) = 0$$

$$x + 1 = 0 \quad x + 6 = 0$$

$$x = -1 \quad x = -6$$

B) bottom up 5

$$m = -1 \quad m = 3$$

quadratic eqn

$$12) n^2 + 3n + 15 = 0$$

$$X$$

$$or$$

$$n = -3, -5$$

Algebra I
Kuta Software LLC. All rights reserved.
U3B SG - Solving Quadratic Functions

Solve each equation by factoring.

1) $(x-1)(5x+8)=0$ 2) $n^2-n-12=0$
 $\boxed{1, -5}$ $\boxed{4, -3}$

3) $7r^2+3r=0$
 $\boxed{-\frac{3}{7}, 0}$

Solve each equation by taking square roots.

4) $a^2 = 25$ 5) $p^2 + 1 = 82$
 $\boxed{5, -5}$ $\boxed{9, -9}$

6) $-4 + 6d^2 = 60$
 $\boxed{1, -1}$

ID: 1

<p>Solve each equation by completing the square.</p> <p>7) $x^2 + 12x + 37 = 8$ [3, -15]</p>	<p>Solve each equation with the quadratic formula.</p> <p>8) $2x^2 - (10 + 11)x = 0$ $\left \frac{5 + \sqrt{3}}{2}, \frac{5 - \sqrt{3}}{2} \right$</p>	<p>Solve each equation by factoring, use any factoring method you choose.</p> <p>9) $k^2 + 7k + 6 = 0$ [-1, -6]</p>	<p>10) $7r^2 - 42r = 0$ [6, 0]</p>
<p>Solve each equation by factoring, use any factoring method you choose.</p> <p>11) $3m^2 - 6m - 9 = 0$ [-1, 3]</p>	<p>12) $n^2 + 8n + 15 = 0$ [-3, -5]</p>		
<p><small>Copyright © Kuta Software LLC. All rights reserved. -2-</small></p>			

March 27, 2019, Wednesday

GSE Algebra I	Solving Quadratics	Notes
Name:	Date:	
A quadratic equation is any equation having the form $ax^2 + bx + c = 0$. A quadratic function forms a graph called a parabola (shaped like a u).		
In this unit, we will solve quadratic equations, meaning we will find the values of x when $y = 0$.		
Solving Quadratics by Graphing		
Solve a Quadratic by Graphing		
To solve a quadratic by graphing is to find where the parabola crosses the x -axis.		
We call these the <u>solutions</u> , <u>roots</u> , <u>zeros</u> , or <u>x-intercepts</u> .		
Example : Find the zeros.		
		Find the solutions. The solutions are the x-intercepts of the parabola, which are $x = -1$ and $x = 2$.
Practice: Identify the solutions of each quadratic graph.		
1.		2.
		3.

Quiz....

You may skip one from the front
& skip one from the back. Please
write "SKIP" on the problems
you do not wish to complete.

Algebra 1
Solving quadratics by graphing
Name _____

EXAMPLE: Use the graphs below to solve the given quadratic equations.

a. $y = -x^2 + 3x + 4$

The graph of $y = -x^2 + 3x + 4$.

b. $x^2 - 6x + 9 = 0$

The graph of $y = x^2 - 6x + 9$.

c. $x^2 + 2x + 4 = 0$

The graph of $y = x^2 + 2x + 4$.

d. $x^2 - 4x - 6 = 0$

The graph of $y = x^2 - 4x - 6$.

SOLUTIONS:

a. Since we are given the graph of $y = -x^2 + 3x + 4$ and we are asked to solve $(x+1)(x+3) = 4$, we need to look for the x -intercepts since the graph of $y = -x^2 + 3x + 4$ intersects the line $y = 0$, i.e., we need to find the x -intercepts of the graph. Since x -intercepts are $(-1, 0)$ and $(4, 0)$, we can conclude that the solution set are $x = -1$ or $x = 4$. So the solution set is $\{-1, 4\}$.

b. Since the graph of $y = x^2 - 6x + 9$ intersects the line $y = 0$ only when $x = 3$, we can conclude that the solution set is $\{3\}$.

c. Since the graph of $y = x^2 + 2x + 4$ never intersects the line $y = 0$ (i.e., the x -axis), we see that $x^2 + 2x + 4 = 0$ never equals to 0, so we can conclude that the equation $x^2 + 2x + 4 = 0$ does not have any real solution. (Remember that we can use the quadratic formula by completing the square to find the complex numbers that are the solution.)

d. Here, we are given the graph of $y = x^2 - 4x - 6$ and we need to find the x when the expression $x^2 - 4x - 6 = 0$. In the given equation $x^2 - 4x - 6 = 0$, which we can do by solving the equation for x :

$$x^2 - 4x - 6 = 0 \quad (\text{add } 6 \text{ to both sides})$$

$$x^2 - 4x = 6$$

We can solve the equation by looking for the x -values where the graph of $y = x^2 - 4x - 6$ intersects the line $y = 0$ (i.e., the x -axis). Thus, the solution set is $\{0, 2\}$.

Algebra 1
Solving quadratics by graphing
Name _____

EXAMPLE 1
Solving quadratics by graphing!

a. $a(x) = (x - 2)^2 - 4$

Vertex: $(2, -4)$
Axis of Symmetry: $x = 2$

b. $b(x) = -2(x - 6)^2 + 4$

Vertex: $(6, 4)$
Axis of Symmetry: $x = 6$

c. $c(x) = -\frac{1}{2}(x + 4)^2 + 2$

Vertex: $(-4, 2)$
Axis of Symmetry: $x = -4$

d. $d(x) = (x + 2)^2$

Vertex: $(-2, 0)$
Axis of Symmetry: $x = -2$

FACTORING
 $x^2 + 5x + 6 = 0$

COMPLETING THE SQUARE
 $x^2 + 8x + 11 = 0$

GRAPHING

QUADRATIC FORMULA

TAKING SQUARE ROOTS

ALWAYS WORKS

SOLVING A QUADRATIC EQUATION

$x^2 = 36$ $y^2 = -40$

+

+

FACTORING
ACB CHART
 $x^2 + 5x + 6 = 0$
 $(x+3)(x+2) = 0$
 $x+3 = 0$ $x+2 = 0$
 $x = -3$ $x = -2$

COMPLETING THE SQUARE

GRAPHING

QUADRATIC FORMULA

TAKING SQUARE ROOTS

ALWAYS WORKS

SOLVING A QUADRATIC EQUATION

$x^2 + 8x + 11 = 0$
 $x^2 + 8x = -11$
 $x^2 + 8x + 16 = -11 + 16$
• Factor:
 $(x+4)^2 = 5$
 $\sqrt{x+4} = \sqrt{5}$
 $x+4 = \pm\sqrt{5}$
 $x = -4 \pm \sqrt{5}$

$x^2 = 36$ $y^2 = -40$

$\sqrt{x^2} = \sqrt{36}$ $\sqrt{y^2} = \sqrt{-40}$
 $x = \pm 6$ $y = 2i\sqrt{10}$

2 REAL SOLUTIONS
ONE REAL SOLUTION
NO REAL SOLUTIONS (TWO COMPLEX)

quiz...

Item 13
Constructed Response

Maria and Jeff collect data on the number of cars that pass through an intersection every Monday morning for 2 months. They record the findings as 57, 63, 63, 71, 56, 67, 76, and 64. They each use different methods to summarize the typical number of cars that pass through the intersection at the specified time and compare their findings. Jeff says that, on average, 79 cars pass through the intersection each Monday morning. Maria disagrees and says that the mean should not be used and uses the median to describe the typical number of cars that pass through the intersection at any given Monday morning.

Part A What is the median value of the data? Write your answer in the space provided.

Part B Explain why the median should be used instead of the mean. Write your answer in the space provided.

Part A _____

Part B _____

